



Nearly 9 out of 10 patients with genotype 1* were able to say...

I am
HEPATITIS Cured

SOVALDI offers hope that you can put hepatitis C (Hep C) behind you. SOVALDI is a prescription medicine used with other antiviral medicines to treat chronic (lasting a long time) hepatitis C genotype 1, 2, 3, and 4 infection in adults. SOVALDI should not be taken alone. It is not known if SOVALDI is safe and effective in children under 18 years of age.

 **DISCOVER SOVALDI**

Leading the way with a breakthrough treatment for Hep C >

 **HOW CAN I GET STARTED?**

Talk to your doctor and see how our team can help >

LEARN ABOUT ANOTHER TREATMENT FOR GENOTYPE 1 >

*In a study of 327 patients who had no prior Hep C treatments and who all received SOVALDI once daily for 12 weeks along with another oral medicine (ribavirin) and a weekly injection (peginterferon alfa), 89% of those with the most common type of Hep C (genotype 1) were cured. Cure means the Hep C virus is not detected in the blood when measured 3 months after treatment is completed.

Important Safety Information

What is the most important information I should know about SOVALDI?

SOVALDI combination therapy with ribavirin or peginterferon alfa and ribavirin can cause birth defects or death of your unborn baby. If you or your sexual partner is pregnant or plans to become pregnant, do not take these medicines.

- You and your sexual partner must use 2 effective forms of birth control and should not become pregnant while being treated with SOVALDI combination therapy and for 6 months after your treatment is over. Talk to your doctor about forms of birth control.
- If you or your partner can become pregnant you must have a pregnancy test before starting treatment with SOVALDI combination therapy, every month while being treated, and for 6 months after your treatment ends.
- Tell your healthcare provider right away if a pregnancy occurs while taking or within 6 months after you stop taking SOVALDI combination therapy. You or your healthcare provider should contact the Ribavirin Pregnancy Registry. If you are also infected with HIV and taking medicines to treat your HIV infection, an Antiretroviral Pregnancy Registry is also available. For contact information, please see [full Prescribing Information](#).

You should not take SOVALDI alone. SOVALDI should be used together with ribavirin or in combination with peginterferon alfa and ribavirin to treat chronic hepatitis C infection.

What should I tell my healthcare provider before taking SOVALDI?

- If you have: liver problems other than hepatitis C infection; had a liver transplant; severe kidney problems or are on dialysis; HIV; any other medical condition; or if you are breastfeeding or plan to breastfeed. You and your healthcare provider should decide if you will take SOVALDI or breastfeed. You should not do both.
- If you take rifampin (Rifadin®, Rifamate®, Rifater®), St. John's wort or a product that contains St. John's wort. Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Other medicines may affect how SOVALDI works.

What are the most common side effects of SOVALDI?

For SOVALDI used in combination with ribavirin they include tiredness and headache and for SOVALDI used in combination with peginterferon alfa and ribavirin they include the additional side effects of nausea, difficulty sleeping, and low red blood cell count.

